



US006410604B1

(12) **United States Patent**
Braue, Jr. et al.

(10) **Patent No.:** **US 6,410,604 B1**
(45) **Date of Patent:** **Jun. 25, 2002**

(54) **ACTIVE TOPICAL SKIN PROTECTANTS
CONTAINING OPAA ENZYMES AND CLECS**

5,607,979 A	3/1997	McCreery	514/759
5,914,436 A	6/1999	Klabunde et al.	588/205
5,990,373 A	11/1999	Klabunde	588/200
6,057,488 A	5/2000	Koper et al.	588/200
6,224,885 B1	5/2001	Jenner	424/401

(75) Inventors: **Ernest H. Braue, Jr.**, Whiteford;
Stephen T. Hobson, Belcamp, both of
MD (US); **Chandrike Govardhan**,
Lexington; **Nazar Khalaf**, Worcester,
both of MA (US)

(73) Assignee: **The United States of America as
represented by the Secretary of the
Army**, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/871,749**

(22) Filed: **Jun. 1, 2001**

Related U.S. Application Data

(60) Provisional application No. 60/209,337, filed on Jun. 2,
2000.

(51) **Int. Cl.**⁷ **A61K 31/02**; A61K 31/08;
A61K 47/00; A61K 7/42

(52) **U.S. Cl.** **514/759**; 424/59; 574/723;
574/772; 574/789; 574/844; 574/845; 574/937;
574/939; 574/944

(58) **Field of Search** 424/59; 514/723,
514/759, 772, 789, 844, 845, 937, 939,
944

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,649,037 A 3/1987 Marsh et al. 423/338

OTHER PUBLICATIONS

Smith, et al., *Jrnl. of the American Acad. of Dermatology*,
Vo. 32, No. 5, part 1, May 1995, pp. 765–776, Sulfur
mustarc: Its continuing threat as a chemical warfare agent,
the cutaneous lesions induced, progress in understanding its
mechanism of action, its long–term health effectgs, and new
developments for protection and therapy.

Arroyo, et al., *Jrnl. of Pharm. and Toxicol. Methods*, vol 33,
No. 2, Apr. 1995, pp. 109–112, *EPR/Spin–Label Technique
as an Analytical Tool for Determining the Resistance of
Reactive Topical Skin Protectants (rTSPs) to the Break-
through of Vesicant Agents.*

Primary Examiner—Shelley A. Dodson

(74) *Attorney, Agent, or Firm*—Elizabeth Arwine

(57) **ABSTRACT**

A topical skin protectant formulation containing a barrier
cream and an active moiety for protecting warfighters and
civilians against all types of harmful chemicals, specifically
chemical warfare agents (CWA's). The topical skin pro-
tectant offers a barrier property and an active moiety that
serves to neutralize chemical warfare agents into less toxic
agents.

34 Claims, 1 Drawing Sheet




FIG. 1

ACTIVE TOPICAL SKIN PROTECTANTS CONTAINING OPAA ENZYMES AND CLECS

PRIORITY INFORMATION

This application claims the benefit of priority of U.S. Provisional Application No. 60/209,337 filed Jun. 2, 2000.

BACKGROUND OF THE INVENTION


1. Field of the Invention

This invention relates to active topical skin protectants. More specifically, the invention relates to an active barrier cream for protection against all types of harmful chemicals, specifically chemical warfare agents (CWAs). The active barrier cream is applied prior to exposure on the skin of persons at risk of exposure to harmful chemicals to provide a protective barrier for the skin. The active barrier cream chemically or physically reacts with harmful chemicals such as CWAs (vesicants and nerve agents) to neutralize these harmful chemicals while the barrier properties of the cream prevent penetration of harmful chemicals through the cream to the skin.

2. Description of Related Art

The concept of applying a topical protectant to vulnerable skin surfaces before entry into a chemical combat arena has been proposed as a protective measure against percutaneous CWA toxicity since the first use of CWAs in World War I. The protectant would be applied to vulnerable skin surfaces prior to entry into a chemical combat area. Topical protectants should augment the protection afforded by the protective over garments and/or redefine the circumstances requiring mission oriented protective posture (MOPP) levels. The rapid action of vesicating agents, also known as blistering agents, such as sulfur mustard (HD) and lewisite (L), require a pre-exposure skin protection system or a contamination avoidance approach that may preclude the percutaneous toxicity of these agents. These approaches also reduce the risk of exposure to organophosphorus (OP) chemical agents (nerve agents) that unlike the vesicating agents, are lethal in droplet amounts.

An organic molecule, S-330, that reacts with CWAs is incorporated into a product and fielded as the M-5 ointment kit at the end of World War II (Formula 1).



However, the unacceptable barrier properties and the undesirable cosmetic properties (i.e. foul odor and sticky texture) caused the cancellation of this product.

Two non-active topical skin protectant (TSP) formulations were developed at the United States Army Medical Research Institute of Chemical Defense (USAMRICD) and were transferred to advanced development following a Milestone Zero (MS0) Review in October 1990. The timeline of the approval of the TSP continued with MSI in 1993, a Investigational New Drug (IND) filed with the FDA in 1994, MSII in 1995, and culminated with New Drug Application

(NDA) approval in February, 2000. Upon approval by the FDA, the TSP was designated Skin Exposure Reduction Paste Against Chemical Warfare Agents (SERPACWA). SERPACWA is a 50:50 (wt/wt) mixture of perfluoropolyether oil (Fomblin® Y25 from Ausimont) and polytetrafluoroethylene (polymist® F5a powder from Ausimont). The formulation described in McCreery U.S. Pat. No. 5,607,979 is directed to a topical skin protectant cream that acts as a barrier to CWAs.

Although SERPACWA extends the protection afforded by MOPP and allows a longer window for decontamination, it does not completely remove the possibility for contamination because the CWA is not neutralized. To avoid contamination of other areas of the battlefield and to preclude the future percutaneous absorption of the CWA, decontamination is still required. Furthermore, despite SERPACWA's excellent protection against GD and HD liquid, its protection against HD vapor is minimal.

To overcome these deficiencies, there is a need for a new TSP that contains an active component. This active Topical Skin Protectant (active TSP) was developed within the following criteria. First, the -active TSP should neutralize CWAs including but not limited to sulfur mustard (HD), soman (GD), and VX. Second, the barrier properties of the TSP should be maintained or increased. Third, the protection against HD vapor should increase. And fourth, the cosmetic characteristics (i.e. odor, texture) of the TSP should be maintained.

This invention meets the above criteria and solves the problems associated with the past TSPs by providing an active topical skin protectant that increases effectiveness of the TSP barrier quality and neutralizes CWAs into less harmful products.

It is therefore, an objective of the present invention to provide an active topical skin protectant that prevents the percutaneous absorption of CWAs and converts these toxic materials into less harmful products.

It is a further objective of the present invention to provide an active topical skin protectant that maintains desirable cosmetic properties making it acceptable to the user. Specifically, the active TSP should not be sticky, without offensive odor, and should be nonirritating to the skin.

It is still a further object of the invention to provide an active topical skin protectant that is practical for field operations. Specifically, the active TSP should have a stable shelf life, not be easily washed off with water, and should not react with insecticides or camouflage paint.

SUMMARY OF THE INVENTION

A topical skin protectant formulation for neutralizing chemical warfare agents into less toxic products comprising: a barrier base cream, and one or more active moieties. The base cream comprises poly(tetrafluoroethylene) (PTFE) resins dispersed in perfluorinated polyether oils (PFPE). The active moieties that have been found to be effective with the base cream are listed in Table 1. The active barrier cream is applied to the skin prior to exposure of persons at risk of exposure to harmful chemicals to provide an active barrier to protect the skin. The active barrier cream chemically or physically reacts with harmful chemicals such as CWA's to neutralize these harmful chemicals while the barrier properties of the cream prevent penetration of harmful chemicals through the cream to the skin.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a flow diagram of the active TSP Decision Tree Network for efficacy evaluation.

DETAILED DESCRIPTION

Candidate Active Moieties

The types of materials that neutralize harmful agents use three main modes of action: oxidation, reduction or hydrolysis.

Operating criteria, however, restricts the selection of the active materials. Thus, the active moiety must not irritate the

skin, react with insecticides or camouflage paints or be unstable. This restriction eliminates many of the most active species. Furthermore, the active moiety must be incorporated into a highly fluorinated environment that is not amenable to many reaction pathways. One such class of compound is enzymes or Cross-Linked Enzyme Crystals (CLECs) that neutralized either nerve or vesicating agents (Table 1).

TABLE 1

LIST OF ACTIVE COMPOUNDS and EXAMPLE FORMULATIONS FOR ACTIVE TOPICAL SKIN PROTECTANTS					
ICD #	Active Moiety	wt % Active	wt % Other	wt % PFPE	Wt % PTFE
3336	OPAA Crystals	1	Polyethylene oxide (300K), glycodeoxycholic acid sodium salt, polyvinyl alcohol (water 7.1%), Tyloxapol, isopropanol, 20 mM HEPES pH 7.2 and Tris(2-carboxyethyl) phosphine HCl	52	40
3337	OPAA Crystals	1	Polyethylene oxide (100K), glycodeoxycholic acid sodium salt, polyvinyl alcohol (water 8.2%), Tyloxapol, isopropanol, 20 mM HEPES pH 7.2 and Tris(2-carboxyethyl) phosphine HCl	52	40
3338	OPAA Crystals	1	glycodeoxycholic acid sodium salt, polyvinyl alcohol (water 7.7%), Tyloxapol, isopropanol, 20 mM HEPES pH 7.2 and Tris(2-carboxyethyl) phosphine HCl	52	40
3339	OPAA Crystals	1	Polyethylene oxide (100K), glycodeoxycholic acid sodium salt (water 7.7%), Tyloxapol, isopropanol, 20 mM HEPES pH 7.2 and Tris(2-carboxyethyl) phosphine HCl	52	40
3340	OPAA CLEC	1	Polyethylene oxide (100K), glycodeoxycholic acid sodium salt (water 23.26%), Tyloxapol, isopropanol, 20 mM HEPES pH 7.2 and Tris(2-carboxyethyl) phosphine HCl	52	40
3622	OPAA	0.95%	Tyloxapol (1.9%), Polyethylene Oxide (300K M. W., 0.18%), Glycodeoxycholic Acid Sodium Salt (0.09%), 10% polyvinyl alcohol (1.85%), 20 mM MOPS (3.69%), Isopropanol (3.69%)	38.1	49.5
3623	OPAA	0.95%	Tyloxapol (1.9%), Polyethylene Oxide (600K M. W., 0.18%), Glycodeoxycholic Acid Sodium Salt (0.09%), 10% polyvinyl alcohol (1.85%), 20 mM MOPS (3.69%), Isopropanol (3.69%)	38.1	49.5
3624	OPAA	0.95%	Tyloxapol (1.9%), Polyethylene Oxide (900K M. W., 0.18%), Glycodeoxycholic Acid Sodium Salt (0.09%), 10% polyvinyl alcohol (1.85%), 20 mM MOPS (3.69%), Isopropanol (3.69%)	38.1	49.5
3625	OPAA	0.95%	Polyethylene Oxide (100K M. W., 0.18%), Glycodeoxycholic Acid Sodium Salt (0.09%), 10% polyvinyl alcohol (1.85%), 20 mM MOPS (3.69%), Isopropanol (0.92%), water (2.77%)	38.1	49.5
3626	OPAA	0.95%	Polyethylene Oxide (100K M. W., 0.19%), Glycodeoxycholic Acid Sodium Salt (0.09%), 10% polyvinyl alcohol (1.88%), 20 mM MOPS (3.76%), Isopropanol (0.94%), water (2.63%), 25% Jeffamine T-4.3 (0.01%)	38.1	49.5
3627	OPAA	0.95%	Polyethylene Oxide (100K M. W., 0.19%), Glycodeoxycholic Acid Sodium Salt (0.10%), 10% polyvinyl alcohol (1.91%), 20 mM MOPS (3.83%), Isopropanol (0.96%), water (2.49%), 25% Jeffamine T-4.3 (0.03%)	38.1	49.5
3201	OPAA Dried Enzyme	10		50	40
3202	OPAA Dried Enzyme	5	H ₂ O (5%), 2853 (5%)	50	35

TABLE 1-continued

LIST OF ACTIVE COMPOUNDS and EXAMPLE FORMULATIONS FOR ACTIVE TOPICAL SKIN PROTECTANTS					
ICD #	Active Moiety	wt %		Wt %	
		Active	Other	PFPE	PTFE
3203	OPAA Dried Enzyme	5	2853 (5%)	50	40
3204	OPAA wet enzyme	10		50	40
3205	OPAA wet enzyme	5	2853 (5%), H2O (5%)	50	35
3206	OPAA wet enzyme	5	2853 (5%)	50	40
3207	OPAA dry enzyme	5	H2O (5%)	50	40
3209	OPAA dry enzyme	5	H2O (5%), 2853 (10%)	45	30
3210	OPAA dry enzyme	5	H2O (15%), 2853 (15%)	35	30

Abbreviations:

PTFE: poly(tetrafluoroethylene) available as F5A powder from Ausimont, Morristown, NJ

PFPE: perfluoropolyether available as FOMBLIM™ Y25 oil from Ausimont, Morristown, NJ

ICD2853: Light PFPE Surfactant, Dupont, Wilmington, DE

OPAA: organophosphorus acid anhydride hydrolase

Percentages are given in weight percents (wt. %).

All active moieties listed above are useful for both liquid and vapor challenges. The amount of each varies with each formulation. The object is to optimize the quantity of active moiety in the base cream without losing the barrier properties of the base cream. The amount of active moiety can vary from about 1–20%. The amount of perfluorinated polyether oil can vary from about 40 to 60%. The amount of poly(tetrafluoroethylene) can vary from about 30 to 50%. One criterion for selection of the active materials is increased efficacy against HD and/or GD vapor. Formulations ICD #3202, 3206, 3336, and 3337 all have significantly (P=0.05) increased protection compared to SERPACWA (ICD 3004) in the penetration cell model against GD. The best candidate compounds listed in Table 1 for GD containing OPAA are ICD3202, ICD3206, ICD3336, and ICD3337.

The enzyme or CLEC must also be incorporated into the TSP matrix without degradation of the barrier properties. The enzymes were incorporated into the cream either as lyophilized solids or aqueous suspensions. Typically they are dispersed into the perfluorinated oil followed by sequential addition of the appropriate amount of F5A PTFE.

SERPACWA (ICD3004) consists of fine particles of poly(tetrafluoroethylene) resin dispersed in a perfluorinated polyether oil. The excellent barrier properties of this high molecular weight polymer formulation are related to the low solubility of most materials in it. Only highly fluorinated solvents like Freon® have been observed to show appreciable solubility. This aprotic non-polar polymer mixture provides a unique medium for active moieties of the invention. Reaction mechanisms that do not involve charged transition states should be favored in this chemical environment.

Base creams formed from about 35–50% fine particulates of certain poly(tetrafluoroethylene) PTFE resins dispersed in perfluorinated polyether oils (PFPE) having viscosities from about 20 cSt to about 500 cSt afford good protection against chemical warfare agents such as HD, L, sulfur mustard/Lewisite mixtures (HL), pinacolyl methylphosphonofluoridate (soman or GD), thickened soman (TGD) and O-ethyl S-(2-diisopropylaminoethyl)methylphosphonothiolate (VX). PTFE and PFPE are available commercially from Ausimont (Morristown, N.J.) and Dupont (Wilmington, Del.).

The base creams used in the invention are suspensions of 35–50% finely divided PTFE having a surface area below about 6 m²/g in a perfluorinated polyether base oil prepared

20

from perfluoropropylene oxide, which has a viscosity between about 20 and about 500 cSt. More preferred compositions comprise from about 35% to about 50% of finely divided PTFE having an average particle size from about 0.1 μm to about 10 μm and a surface area below about 4 m²/g in a perfluorinated polyether base oil from 40% to 60% having a viscosity between about 20 and about 500 cSt.

Suitable perfluorinated polyether oils are Fomblin® HC- and Y-oils (Ausmont) and Krytox® oils (Dupont). The Fomblin® oils are mixtures of linear polymers based on perfluoropropylene oxide having the following chain structure: CF₃—[(OCF(CF₃)CF₂)_n—(OCF₂)_m]OCF₃. The Krytox® oils are mixtures of linear polymers also based on perfluoropropylene oxide and have the chemical structure F—[(CF(CF₃)CF₂O)]_mCF₂CF₃. Fomblin® Z oils having the formula: CF₃—[(OCF₂CF₂)_n—(OCF₂)_m]—OCF₃, may also be useful in the practice of the invention. The indices n and m indicate the average number of repeating polymeric subunits in the oil molecules. The oils may have a viscosity of about 20 cSt to about 500 cSt or more. The creams were generally prepared according to U.S. Pat. No. 5,607,979, incorporated herein in its entirety.

Other additives to the base cream are water and surfactant and other chemicals necessary to maintain OPAA activity (see Table 1). The surfactant facilitates the mixing of the water with the base cream. An example of a typical surfactant is perfluoropolyalkylether (Krytox® CAS #60164-51-4 from Dupont). Additional materials may also be incorporated as long as they do not reduce effectiveness of the topical protectant, such as stabilizers, camouflage paints, and sunscreens.

A further understanding of the composition of the topical protectant of the invention can be obtained by reference to certain specific example formulations set forth in Table 1. These examples are provided herein for purposes of illustration only and are not intended to be limiting. Many active moieties require the presence of water as a reagent for the hydrolysis of HD and GD. The active moieties that react by a hydrolysis mechanism require the presence of water. When the topical protectant is applied to the skin of a user, moisture in the form of perspiration may also aid in the hydrolysis of HD and GD. The addition of perfluorinated polyether surfactants to the base cream facilitates the addition of water.

Temperature and mixing shear should be monitored to maintain the base cream at the desired consistency and

quality. The active TSPs are typically prepared at ambient temperature using mechanical mixing. Mixing times of 10–20 minutes are usually sufficient for dispersal of enzymes into the SERPACWA matrix. A typical procedure for the preparation of an active aTSP with enzymes is presented below:

In a polypropylene container is added the appropriate amount of enzyme (1–3% by weight) and Y25 (50–55% by weight) perfluorinated oil. The suspension is mixed with a mechanical stirrer at ambient temperature for 5 to 15 minutes. To the suspension is added F5A poly (tetrafluoroethylene) in three portions with vigorous mechanical stirring of 5 to 10 minutes between each addition. After final addition the container is tightly capped and sealed with Parafilm®.

Multilayer Approach

Although an active TSP can be generally the application of a powder that is a POM/RNP sprinkled on the skin, or an active moiety in a base cream wherein the cream is spread on the skin, a multilayering approach can also be used. The multilayer approach would be to use the active TSP as the first layer and a solid active moiety powder as the second layer. The second layer would be a thin coating of the solid active moiety powder sprinkled over the active TSP cream. This approach would provide a concentrated decontamination material at the surface of the barrier cream, which would accelerate the neutralization process of CWA's coming in contact with the surface. In the alternative, one or more solid active moiety powders can be applied first followed by an application of the active TSP.

Testing

Evaluation of formulations was conducted with a decision tree network (DTN) that describes the path that active TSPs follow during evaluation (FIG. 1).

The DTN is divided into two pathways: one for vesicants and the other for nerve agents. Within these pathways there are three blocks each with a decision point. The first block consists of a series of three mechanical (in vitro) modules used to determine the initial efficacy of candidate formulations and to eliminate non-effective candidates before animal testing, the second block consists of in vivo modules and the third block consists of an advanced animal module to determine the influence of time, water and interactions with other products.

The M8 paper test is used to evaluate the barrier resistance of liquid CWA challenges, including HD, pinacolyl methylphosphonofluoridate (soman, GD), and O-ethyl S-(2-diisopropylaminoethyl) methylphosphonothioate (VX). In this test a 0.15 mm layer of active TSP is placed over a well-defined area of M8 chemical detection paper and challenged with an 8 μ l droplet of CWA. When agent penetrates the active TSP barrier and reaches the M8 paper, a colored spot develops on the paper. The test assemblies are observed for 6 hr and the breakthrough time is reported for each sample. A total of nine replicates are run for each test, and a standard reference compound is included each day for quality control.

The penetration cell test is used to evaluate the barrier properties against both liquid and vapor CWA challenges (Braue, E. H. Jr. *Journal of applied Toxicology*, 1999, 19(S), S47–S53). In this test the lower half of a Reifenrath diffusion cell (Reifenrath Consulting and Research, Richmond, Calif.) is used. A 0.15 mm thick layer of active TSP is supported by nitrocellulose paper on top of the cell. The active TSP layer

is challenged with a 10- μ l liquid droplet of HD or an 8 μ l droplet of GD, or a saturated vapor cup of HD or GD. Breakthrough of CWA into the lower chamber of the diffusion cell is monitored using a miniature continuous air monitoring system (MINICAMS, CMS Research, Birmingham, Ala.). This system has been automated to allow continuous monitoring of five cells in a 40-min cycle. The test runs for 20 hr and the accumulated amounts of agent that break through the active TSP barrier are calculated. From these data, we obtained two values: the cumulative amount of CWA that penetrates through the active TSP, and the time at which a “breakthrough” occurs. We defined “breakthrough” values at the minimum amount of HD (1000 ng) and GD (1000 ng) that results in a physiological response. Minimal amount of HD for vesication=1000 ng. See F. R. Sidell, J. S. Urbanetti, W. J. Smith, and C. G. Hurst in *Textbook of Military Medicine, Medical Aspects of Chemical and Biological Warfare*, edited by F. R. Sidell, E. T. Takafuji, and D. R. Franz (Office of the Surgeon General at TMM Publications, Washington, D.C. 1997) p 201. LD₅₀ for soman (GD)=350 mg/70 kg man. See F. R. Sidell in *Textbook of Military Medicine, Medical Aspects of Chemical and Biological Warfare*, edited by F. R. Sidell, E. T. Takafuji, and D. R. Franz (Office of the Surgeon General at TMM Publications, Washington, D.C. 1997) p 141. These two values allow us to rank the active TSP formulations and to select the appropriate component for advanced development.

The proof-of-neutralization test is used to verify that active TSP formulations actually neutralize CWAs into less toxic materials. This test uses the headspace solid phase microextraction (HS-SPME) technique for the collection of CWAs. Samples collected on the extraction filament are analyzed by gas chromatography/mass spectroscopy. 100 mg of active TSP formulation are challenged with 0.1 μ l of neat CWA (HD, GD, or VX) in a small vial. The headspace above the mixture is sampled periodically to determine the amount of CWA remaining in the flask. Efficacy is determined by the % loss of CWA. Other analytical techniques such as Nuclear Magnetic Resonance (NMR) and Fourier-Transform Infrared Spectrometry (FTIR) have also been used in this module.

Formulations that pass this initial set of screens are moved into the second phase of testing using animal models. The weanling pig test for HD vapor evaluates a 0.10 to 0.20 mm thick layer of active TSP spread on the depilated dorsa. The standard saturated vapor cup is used for a 15–60 min challenge. The effectiveness of the active TSP is determined by measuring the degree of erythema that developed on the skin exposure site. Erythema is measured objectively using a reflectance colorimeter (see Braue, E. H. Jr. *Journal of Applied Toxicology*, 1999, 19(S), S47–S53).

The rabbit lesion area ratio (LAR) test is used to evaluate a challenge by HD liquid. In this test, a 0.10 mm layer of active TSP spread on the clipped dorsa is challenged with 1.0 μ l of liquid HD. The effectiveness of the active TSP is determined by measuring the lesion areas of protected and non-protected sites.


The rabbit acetylcholinesterase (ACHE) inhibition test is performed by applying a 0.10 mm thick layer of active TSP on the clipped dorsa of rabbit followed by a fixed dose of GD (1 LD₅₀), TGD (1 LD₅₀), or VX (20 LD₅₀). The effectiveness of the active TSP is determined by lethality and also by measuring the erythrocyte acetylcholinesterase activity 0.5, 1, 2, and 24 hr following exposure.

Candidate formulations that pass the in vivo test modules move into advanced animal testing. These tests are similar to

the initial animal tests with the addition of stresses for wear-time and washing with water. Interactions with other products that a soldier might use are also evaluated. These products include camouflage paints, sunscreens and insecticides.

Results

With the aforementioned enzymes and CLECs, the expected neutralization mechanism is hydrolysis (Scheme 1).



Since the OPAA and OPH enzymes are specific for organophosphate hydrolysis, it is expected that only the hydrolysis of GD will be accelerated and the hydrolysis of HD will be due to the residual water or other additives (see Table 1). The increase in protection for the active TSPs containing enzymes and CLECs was most evident against GD vapor. This is shown by the increase in the time needed for 1000 ng of GD vapor to penetrate the active TSP as compared to SERPACWA (ICD 3004) (Chart 1). Formulations ICD #3202, 3206, 3336, and 3337 all have significantly increased protection compared to SERPACWA (ICD 3004) in the penetration cell model against GD.

barrier base cream; and one or more (OPAA) organophosphorus acid anhydride hydrolase crystals or cross linked enzyme crystals (CLECs) as an active moiety.


2. A topical skin protectant formulation for neutralizing chemical warfare agents into less toxic products comprising a base cream, wherein the base cream comprises poly(tetrafluoroethylene) resins dispersed in perfluorinated polyether oils; and one or more (OPAA) organophosphorus acid anhydride hydrolase crystals or cross linked enzyme crystals (CLECs) as an active moiety.

3. The topical protectant formulation of claim 2, wherein said cross-linked enzyme crystals comprise organophosphorus acid anhydride hydrolase or organophosphorus acid hydrolase.

4. Topical skin protectant system comprising:

- a) a topical skin protectant formulation for neutralizing chemical warfare agents into less toxic products comprising a barrier cream and one or more active moieties; and
- b) a second formulation for applying a thin solid active moiety powder coating comprising one or more active moieties selected from the group consisting of OPAA

Chart 1
Time for 1000 ng to penetrate activate TSPs containing enzymes and CLECS



60

Having now fully described the invention, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the invention as set forth herein.

What is claimed is:

1. A topical skin protest formulation for neutralizing chemical warfare agents into less toxic products comprising:

crystals, OPAA CLEC, Organophosphorus acid hydrolase, Light PFPE Surfactant, Medium PFPE Surfactant, and Heavy PFPE surfactant.

5. A method of protecting a user against chemical warfare agents comprising:

applying a topical skin protectant formulation for neutralizing chemical warfare agents into less toxic prod-

ucts comprising: (a) a barrier cream and (b) one or more active moieties, said one or more active moieties selected from the group consisting of: OPAA crystals, OPAA CLEC, Organophosphorus acid hydrolase, Light PFPE Surfactant, Medium PFPE Surfactant, and Heavy PFPE surfactant.

6. A method of making a topical skin protectant formulation comprising:

mixing one or more active moiety selected from the group consisting of OPAA crystals, OPAA CLEC, Organophosphorus acid hydrolase, Light PFPE Surfactant, Medium PFPE Surfactant, and Heavy PFPE surfactant with a barrier cream comprising poly(tetrafluoroethylene) resins dispersed in perfluorinated polyether oils.

7. A topical skin protectant formulation for neutralizing chemical warfare agents into less toxic products comprising:

(a) a barrier base cream, said barrier base cream comprising poly(tetrafluoroethylene), resins dispersed in perfluorinated polyether oils; and

(b) one or more active moieties, wherein said one or more active moieties is OPAA crystals or cross linked enzyme crystals.

8. The topical skin protectant formulation of claim 7, further comprising one or more additives.

9. The topical skin protectant formulation of claim 8, wherein said additives comprise one or more of water, stabilizers, camouflage paints, and sunscreens.

10. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

(a) about 1 wt. % organophosphorus acid anhydride hydrolase crystals,

(b) about 52 wt. % perfluoropolyether,

(c) about 40 wt % poly(tetrafluoroethylene),

(d) polyethylene oxide (300K),

(e) glycodeoxycholic acid sodium salt,

(f) polyvinyl alcohol water,

(g) tyloxapol,

(h) isopropanol,

(i) 20 mM HEPES pH 7.2 and

(j) tris (2-carboxyethyl) phosphine HCl.

11. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 1 wt. % organophosphorus acid anhydride hydrolase crystals,

b) about 52 wt. % perfluoropolyether,

c) about 40 wt. % poly(tetrafluoroethylene),

d) polyethylene oxide (100K),

e) glycodeoxycholic acid sodium salt,

f) polyvinyl alcohol (water 8.2%),

g) tyloxapol,

h) isopropanol,

i) 20 mM HEPES pH 7.2, and

j) tris(2-carboxyethyl) phosphine HCl.

12. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 1 wt. % organophosphorus acid anhydride hydrolase crystals,

b) about 52 wt. % perfluoropolyether,

c) about 40 wt. % poly(tetrafluoroethylene)

d) glycodeoxycholic acid sodium salt,

e) polyvinyl alcohol (water 7.7%),

f) tyloxapol,

g) isopropanol,

h) 20 mM HEPES pH 7.2, and

i) tris(2-carboxyethyl) phosphine HCl.

13. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 1 wt. % organophosphorus acid anhydride hydrolase crystals

b) about 52 wt. % perfluoropolyether,

c) about 40 wt. % poly(tetrafluoroethylene),

d) polyethylene oxide (100K),

e) glycodeoxycholic acid sodium salt (water 7.7%),

f) tyloxapol,

g) isopropanol,

h) 20 mM HEPES pH 7.2, and

i) Tris(2-carboxyethyl) phosphine HCl.

14. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 1 wt. % organophosphorus acid anhydride hydrolase or cross linked enzyme crystals,

b) about 52 wt. % perfluoropolyether,

c) about 40 wt. % poly(tetrafluoroethylene),

d) polyethylene oxide (100K),

e) glycodeoxycholic acid sodium salt (water 23.26%),

f) tyloxapol,

g) isopropanol,

h) 20 mM HEPES pH 7.2, and

i) Tris(2-carboxyethyl) phosphine HCl.

15. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 0.95 wt. % organophosphorus acid anhydride hydrolase,

b) about 38.1 wt. % perfluoropolyether,

c) about 49.5 wt. % poly(tetrafluoroethylene),

d) tyloxapol (1.9%),

e) polyethylene oxide (300K M. W., 0.18%),

f) glycodeoxycholic acid sodium salt (0.09%),

g) 10 wt. % polyvinyl alcohol (1.85%),

h) 20 mM MOPS (3.69%), and

a) isopropanol (3.69%).

16. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 0.95 wt. % organophosphorus acid anhydride hydrolase,

b) 38.10 wt. % perfluoropolyether,

c) 49.50 wt. % poly(tetrafluoroethylene),

i) tyloxapol (1.9%),

b) polyethylene oxide (600K M. W., 0.18%),

c) glycodeoxycholic acid sodium salt (0.09%),

d) 10 wt. % polyvinyl alcohol (1.85%),

e) 20 mM MOPS (3.69%), and

f) isopropanol (3.69%).

17. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

a) about 0.95 wt. % organophosphorus acid anhydride hydrolase,

b) about 38.1 wt. % perfluoropolyether,

c) about 49.5 wt. % poly(tetrafluoroethylene),

d) tyloxapol (1.9%),

13

- e) polyethylene oxide (900K M. W., 0.18%),
- f) glycodeoxycholic acid sodium salt (0.09%),
- g) 10 wt. % polyvinyl alcohol (1.85%),
- h) 20 mM MOPS (3.69%), and
- i) Isopropanol (3.69%).

18. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 0.95 wt. % organophosphorus acid anhydride hydrolase,
- b) about 38.1 wt. % perfluoropolyether,
- c) about 49.5 wt. % poly(tetrafluoroethylene),
- d) polyethylene oxide (100K M. W., 0.18%),
- e) glycodeoxycholic acid sodium salt (0.09%),
- f) 10 wt. % polyvinyl alcohol (1.85%),
- g) 20mM MOPS (3.69%),
- h) isopropanol (0.92%), and
- i) water (2.77%).

19. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 0.95 wt. % organophosphorus acid anhydride hydrolase,
- b) about 38.1 wt. % perfluoropolyether,
- c) about 49.5 wt. % poly(tetrafluoroethylene),
- d) polyethylene oxide (100K M. W., 0.19%),
- e) glycodeoxycholic acid sodium salt (0.09%),
- f) 10 wt. % polyvinyl alcohol (1.88%),
- g) 20 mM MOPS (3.76%),
- h) isopropanol (0.94%),
- i) water (2.63%), and
- j) 25% Jeffamine T-4.3 (0.01%).

20. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 0.95 wt. % organophosphorus acid anhydride hydrolase,
- b) about 38.1 wt. % perfluoropolyether,
- c) about 49.5 wt. % poly(tetrafluoroethylene),
- d) polyethylene oxide (100K M. W., 0.19%),
- e) glycodeoxycholic acid sodium salt (0.10%),
- f) 10 wt. % polyvinyl alcohol (1.91%),
- g) 20 mM MOPS (3.83%),
- h) isopropanol (0.96%),
- i) water (2.49%), and
- j) 25% Jeffamine T-4.3 (0.03%).

21. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 10 wt. % organophosphorus acid anhydride hydrolase dried enzyme,
- b) about 50 wt. % perfluoropolyether, and
- c) about 40 wt. % poly(tetrafluoroethylene).

22. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 5 wt. % organophosphorus acid anhydride hydrolase dried enzyme,
- b) about 50 wt. % perfluoropolyether,
- c) about 35 wt. % poly(tetrafluoroethylene),
- d) about 5 wt. % water, and
- e) about 5 wt. % Light PFPE surfactant.

23. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

14

- a) about 5 wt. % organophosphorus acid anhydride hydrolase dried enzyme,
- b) about 50 wt. % perfluoropolyether,
- c) about 40 wt. % poly(tetrafluoroethylene), and
- d) about 5 wt. % Light PFPE surfactant.

24. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 10 wt. % organophosphorus acid anhydride hydrolase wet enzyme,
- b) about 50 wt. % perfluoropolyether, and
- c) about 40wt. % poly(tetrafluoroethylene).

25. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 5 wt. % organophosphorus acid anhydride hydrolase wet enzyme,
- b) about 50 wt. % perfluoropolyether,
- c) about 35 wt. % poly(tetrafluoroethylene),
- d) about 5 wt. % Light PFPE surfactant, and
- e) about 5 wt. % water.

26. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 5 wt. % organophosphorus acid anhydride hydrolase wet enzyme,
- b) about 50 wt. % perfluoropolyether,
- c) about 40 wt. % poly(tetrafluoroethylene), and
- d) about 5 wt. % Light PFPE surfactant.

27. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 5 wt. % organophosphorus acid anhydride hydrolase dried enzyme,
- b) about 50 wt. % perfluoropolyether,
- c) about 40 wt. % poly(tetrafluoroethylene), and
- d) about 5 wt. % water.

28. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 5 wt. % organophosphorus acid anhydride hydrolase dried enzyme,
- b) about 45 wt. % perfluoropolyether,
- c) about 30wt. % poly(tetrafluoroethylene),
- d) about 10 wt. % water, and
- e) about 10 wt. % Light PFPE surfactant.

29. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- a) about 5 wt. % organophosphorus acid anhydride hydrolase dried enzyme,
- b) about 35 wt. % perfluoropolyether,
- c) about 30 wt. % poly(tetrafluoroethylene),
- d) about 15 wt. % water, and
- e) about 15 wt. % Light PFPE surfactant.

30. A topical skin protectant formulation of claim 8, wherein said formulation comprises:

- (a) about 1–20 wt. % organophosphorus acid anhydride hydrolase crystals or CLECs,
- (b) about 40–60 wt. % perfluoropolyether, and
- (c) about 30–50 wt. % poly(tetrafluoroethylene).

31. The topical skin protectant formulation of claim 30, further comprising one or more additives selected from the group consisting of polyethylene oxide (300K), glycodeoxycholic acid sodium salt, polyvinyl alcohol, water, tyloxapol, isopropanol, 20 mM HEPES pH 7.2, light surfactant, 20 mM MOPS, Jeffamine T-4.3, and Tris (2-carboxyethyl) phosphine HCl.

15

32. A topical skin protectant formulation for neutralizing chemical warfare agents into less toxic products comprising one or more active moieties, wherein said active moiety is (OPAA) organophosphorus acid anhydride hydrolase crystals or cross linked enzyme crystals (CLECs).

33. The topical skin protectant formulation of claim **32**, further comprising a base cream.

16

34. The topical skin protectant formulation of claim **2**, wherein the active moiety is present in an amount of about 1–20%, the perfluorinated polyether oil is present in an amount of about 40 to 60%, and the poly(tetrafluoroethylene) is present in an amount of about 30 to 50%.

* * * * *